

Loss of least-loaded class in asexual populations due to drift and epistasis

Kavita Jain

Theoretical Sciences Unit and Evolutionary and Organismal Biology Unit,
Jawaharlal Nehru Centre for Advanced Scientific Research,
Jakkur P.O., Bangalore 560064, India

May 30, 2008

Running head: Loss of least-loaded class

Keywords: asexual evolution, Muller's ratchet, epistasis

Corresponding author:

Kavita Jain,

Theoretical Sciences Unit and Evolutionary and Organismal Biology Unit,
Jawaharlal Nehru Centre for Advanced Scientific Research,
Jakkur P.O., Bangalore 560064, India.

jain@jncasr.ac.in

Abstract: We consider the dynamics of a non-recombining haploid population of finite size which accumulates deleterious mutations irreversibly. This ratchet like process occurs at a finite speed in the absence of epistasis, but it has been suggested that synergistic epistasis can halt the ratchet. Using a diffusion theory, we find explicit analytical expressions for the typical time between successive clicks of the ratchet for both non-epistatic and epistatic fitness functions. Our calculations show that the inter-click time is of a scaling form which in the absence of epistasis gives a speed that is determined by size of the least-loaded class and the selection coefficient. With synergistic interactions, the ratchet speed is found to approach zero rapidly for arbitrary epistasis. Our analytical results are in good agreement with the numerical simulations.

In an asexual population of size N , even the fittest individuals can be lost by stochastic fluctuations arising due to the finiteness of the population size. If the individual's genome is long enough that the back mutations can be ignored and recombination is absent, the minimum number of deleterious mutations (least-loaded class) in a finite population increases irreversibly (MULLER, 1964; FELSENSTEIN, 1974). For this reason, this process has been invoked as a potential cause for the evolution of sex and recombination (JUDSON and NORMARK, 1996; HURST and PECK, 1996; BARTON and CHARLESWORTH, 1998), degeneration of nonrecombining parts like Y chromosome (CHARLESWORTH, 1978) and mitochondrial DNA (LOEWE, 2006) of sexually reproducing organisms and extinction of obligately asexual populations by mutational meltdown (GABRIEL *et al.*, 1993).

Due to the irreversible accumulation of deleterious mutations, the process described above acts like a ratchet each click of which corresponds to the loss of the least-loaded class. In the simplest model known as the Muller's ratchet, the ratchet clicks at a constant rate which depends on the population size N , mutation rate U and selection coefficient s . The ratchet speed is also known to depend on other biologically relevant factors such as recombination rate (BELL, 1988; CHARLESWORTH *et al.*, 1993), epistatic interactions (KONDRASHOV, 1994; BUTCHER, 1995; SCHULTZ and LYNCH, 1997), fraction and selection coefficient of favorable mutations (WOODCOCK and HIGGS, 1996; BACHTROG and GORDO, 2004) and spatial structure of the population (COMBADÃO *et al.*, 2007). Although much numerical data for the ratchet speed is available, very few analytical results are known.

As it is desirable to stop or at least slow down the ratchet, several mechanisms with this objective have been proposed (BELL, 1988; WAGNER and GABRIEL, 1990; CHARLESWORTH *et al.*, 1993). One such proposal is to include epistatic interactions in the genome fitness (CHARLESWORTH *et al.*, 1993; KONDRASHOV, 1994). It has been observed experimentally that the gene loci do not always contribute independently to the genome fitness (WOLF *et al.*, 2000; DE VISSER and ELENA, 2007) and the effect of two deleterious mutations can be better (antagonistic) or worse (synergistic) than were they to act independently. For Muller's ratchet operating under epistatic selection, it was noted using numerical simulations that "sufficiently strong synergistic epistasis can effectively halt the action of Muller's ratchet" (KONDRASHOV, 1994). However it was not shown how the ratchet speed approaches zero asymptotically and how much epistatic interaction is required to halt the ratchet.

In this article, besides the classic Muller’s ratchet that assumes haploid asexual population evolving on a fitness landscape in which each gene locus contributes independently to the fitness of the genome, we also study Kondrashov’s model which considers fitness functions with epistatic interactions. We assume that an individual with k mutations has a fitness

$$W(k) = (1 - s)^{k^\alpha} \quad (1)$$

where s is the selection coefficient. For $\alpha = 1$, the epistatic interactions are absent while $\alpha > 1$ corresponds to synergistically epistatic fitness. Our main purpose is to obtain explicit analytical expressions for the typical time $\bar{T}_J$ elapsed between the $(J - 1)$ th and J th click of the ratchet in these models. If the time spent between any two successive clicks is found to be constant then the ratchet turns with a finite speed $1/\bar{T}$, while it is said to be halted at large times if the inter-click time increases with the number of accumulated mutations.

In the past, Muller’s ratchet has been investigated using a diffusion approximation which assumes that the population n_0 of the least loaded class is large and applies to slowly clicking ratchet (STEPHAN *et al.*, 1993; GORDO and CHARLESWORTH, 2000a; STEPHAN and KIM, 2002). The opposite situation of small n_0 and high ratchet rate has been described by a moments method (GABRIEL *et al.*, 1993; GESSLER, 1995; HIGGS and WOODCOCK, 1995; PRÜGEL-BENNETT, 1997; ROUZINE *et al.*, 2003). In this article, we adopt the first method that works in the parameter range for which $n_0 \gg 1$ and the ratchet clicks are slow enough that the population can equilibrate between successive clicks. This requires the knowledge of the steady state properties of an infinitely large population which are known exactly for the Muller’s ratchet (KIMURA and MARUYAMA, 1966; HIGGS, 1994) but have been studied numerically for epistatic case (KONDRASHOV, 1994). After defining the models in the following section, we solve the deterministic quasispecies equation in steady state for $\alpha > 1$ and show that the population frequency of the class with minimum number J of mutations increases with J . These deterministic results are then used to find an expression for the typical time $\bar{T}_J$ in terms of a double integral over the frequency of the least-loaded class which have been evaluated numerically for $\alpha = 1$ (STEPHAN *et al.*, 1993; GORDO and CHARLESWORTH, 2000a). Here we estimate these integrals analytically and find that for a broad range of parameters, the average inter-click time is of a scaling form (29) for any $\alpha \geq 1$. For $\alpha = 1$, it is shown that the ratchet speed is a function of the number $n_0 = Ne^{-U/s}$ in the least-loaded class and the selection coefficient s , and

not n_0 alone as assumed in previous studies (HAIGH, 1978). With epistatic interactions, the time $\overline{T}_J$ is found to increase faster than any power law with J for any $\alpha > 1$. Thus, an arbitrarily small amount of epistasis is sufficient to halt the ratchet with the ratchet speed approaching zero as $\sim 1/t$ for large time t .

MODELS

We consider a haploid asexual population evolving via mutation-selection dynamics starting with an initial condition in which all the individuals in the population have zero mutations. The genome length is assumed to be infinite so that back mutations can be ignored. If the population has a finite size N , it evolves stochastically following the discrete time Wright-Fisher dynamics. An offspring in generation $t + 1$ chooses a parent in the previous generation with a probability proportional to the fitness of the parent. Then the probability $P(n)$ that a parent p carrying k mutations and with fitness $W(k; p)$ has n descendants in one generation is given by

$$P(n) = \binom{N}{n} \left(\frac{W(k; p)}{N\langle W \rangle} \right)^n \left(1 - \frac{W(k; p)}{N\langle W \rangle} \right)^{N-n} \quad (2)$$

where $\langle W \rangle = \sum_{k=0}^{\infty} W(k)X(k, t)$ is the average fitness of the finite population in generation t . Here we have defined $X(k, t)$ as the fraction of population with k mutations in a single sampling of Wright-Fisher process. From the above equation, it follows that the average number of offspring produced in one generation is proportional to the parent's fitness and the relative variance in offspring number decays as $1/N$. This fact will be useful in defining the diffusion coefficient (24) within diffusion approximation discussed in later section. Following replication, mutations are introduced where the number of new mutations is a random variable chosen from a Poisson distribution with mean U . In the simulations, the above process was implemented but the order of mutation and selection was reversed. An individual picked randomly from the population at time t was first mutated and the resulting mutant was allowed to survive at $t + 1$ with a probability equal to its fitness. This process was repeated until the generation $t + 1$ has N members and the population fraction $X(k, t + 1)$ was recorded. It is useful to define $X_J(k, t) = X(J + k, t)$ where J is the minimum number of mutations in the population at time t so that $X_J(k, t) = 0$ for $k < 0$. If $X_J(0, \tau_J)$ becomes zero, the least-loaded class J is lost and the ratchet has clicked at time τ_J .

The ratchet effect due to which the least-loaded class is lost is essentially a stochastic problem arising due to the finite number N of individuals in the population. However as we shall describe later, the population fluctuates close to the deterministic frequency between two clicks of the ratchet. For this reason, we also study the problem of an infinite population for which the density fluctuations vanish and the average population fraction $\mathcal{X}(k, t)$ with k mutations at time t obeys a deterministic quasispecies equation (EIGEN, 1971; JAIN and KRUG, 2007). Similar to the finite population problem, we define $\mathcal{X}_J(k, t) = \mathcal{X}(J + k, t)$ where $\mathcal{X}_J(k, t) = 0$ for $k < 0$. Then neglecting the back mutations for a genome of infinite length (HIGGS, 1994), the fraction $\mathcal{X}_J(k, t)$ evolves according to the following difference equation,

$$\mathcal{X}_J(j, t + 1) = \frac{1}{\mathcal{W}_J(t)} \sum_{k=0}^j e^{-U} \frac{U^{j-k}}{(j-k)!} W(J+k) \mathcal{X}_J(k, t) . \quad (3)$$

In this equation, the population fraction with k mutations replicates with fitness $W(k)$ and accumulates further mutations which are Poisson distributed with a mean U . The average fitness $\mathcal{W}_J(t) = \sum_{k=0}^{\infty} W(J+k) \mathcal{X}_J(k, t)$ in the denominator ensures that the number density is conserved.

STEADY STATE OF THE QUASISPECIES MODEL

In this section, we calculate the steady state population frequency $\mathcal{X}_J(k)$ in the error class $J + k$. Unlike for the multiplicative fitness case, the frequency $\mathcal{X}_J(k)$ depends on J for the epistatic fitness function (HAIGH, 1978). In particular, the fraction $\mathcal{X}_J(0)$ (later abbreviated as $\mathcal{X}_J$) in the least-loaded class is expected to increase with J for $\alpha > 1$ and decrease for $\alpha < 1$. This can be explained by a simple argument which has also been used to understand the error threshold phenomenon (EIGEN, 1971; JAIN and KRUG, 2007) in which the fittest genomic sequence can get lost beyond a critical mutation rate in populations evolving on epistatic fitness landscapes (WIEHE, 1997). Consider the ratio $W(J+k)/W(J) \sim (1-s)^{\alpha k J^{\alpha-1}}$ for $J \gg k$. For synergistic interactions, the error class $J+k$ in the neighborhood of the least-loaded class has a fitness much worse than the fitness of class J rendering selection effective in localising population in the class with J mutations. With increasing J , the selection pressure increases further. Thus we may expect the population frequency $\mathcal{X}_J(k)$ to peak around J and $\mathcal{X}_J$ to increase with J . On the other hand, in case of antagonistic epistasis ($\alpha < 1$), the fitness

landscape is nearly neutral at large J so that the least-loaded sequence can be lost even in the deterministic limit (finite error threshold) (WIEHE, 1997).

In the steady state, the quasispecies equation (3) reduces to

$$\mathcal{X}_J(j) = \frac{1}{\mathcal{W}_J} \sum_{k=0}^j e^{-U} \frac{U^{j-k}}{(j-k)!} W(J+k) \mathcal{X}_J(k) \quad (4)$$

where $\mathcal{W}_J$ is the average fitness in the steady state when the least-loaded class is J . The equation for $j = 0$ immediately shows that (KIMURA and MARUYAMA, 1966; HAIGH, 1978)

$$\mathcal{W}_J = W(J) e^{-U} . \quad (5)$$

For $j = 1$ in (4), we have

$$\mathcal{X}_J(1) = \frac{UW(J)\mathcal{X}_J}{W(J) - W(J+1)} .$$

Plugging this expression in the equation for $j = 2$, after some algebra we obtain

$$\mathcal{X}_J(2) = \frac{U^2W(J)\mathcal{X}_J}{W(J) - W(J+2)} \left[\frac{1}{2} + \frac{W(J+1)}{W(J) - W(J+1)} \right] .$$

Similarly, the fraction in the error class $J+3$ is

$$\begin{aligned} \mathcal{X}_J(3) &= \frac{U^3W(J)\mathcal{X}_J}{W(J) - W(J+3)} \left[\frac{1}{3!} + \frac{W(J+1)}{2!(W(J) - W(J+1))} \right. \\ &\quad \left. + \frac{W(J+2)}{W(J) - W(J+2)} \left(\frac{1}{2!} + \frac{W(J+1)}{W(J) - W(J+1)} \right) \right] . \end{aligned}$$

From the expressions for $\mathcal{X}_J(k)$ for $k = 2, 3$ shown above, it is clear that in the weak selection limit $s \rightarrow 0$, the leading order contribution to $\mathcal{X}_J(k)$ comes from the last term. In general, we can write

$$\begin{aligned} \mathcal{X}_J(k) &\approx \frac{U^k W(J) \mathcal{X}_J}{W(J+k)} \prod_{m=1}^k \frac{W(J+m)}{W(J) - W(J+m)} \\ &\approx \left(\frac{U}{s} \right)^k \mathcal{X}_J \prod_{m=1}^k \frac{1}{(J+m)^\alpha - J^\alpha} \end{aligned} \quad (6)$$

where the population $\mathcal{X}_J$ in the least-loaded class is determined using the normalisation condition $\sum_{k=0}^{\infty} \mathcal{X}_J(k) = 1$.

Using the preceding equation for the multiplicative fitness function, we obtain the well known result (KIMURA and MARUYAMA, 1966)

$$\mathcal{X}_J(k) = \left(\frac{U}{s}\right)^k \frac{\mathcal{X}_J}{k!} = \left(\frac{U}{s}\right)^k \frac{e^{-U/s}}{k!} . \quad (7)$$

The fraction $\mathcal{X}_J(k)$ for all k is seen to be independent of J (HAIGH, 1978). For synergistically epistatic fitness landscape with $\alpha = 2$ in fitness function (1), we have

$$\mathcal{X}_J(k) \approx \left(\frac{U}{s}\right)^k \mathcal{X}_J \prod_{m=1}^k \frac{1}{2Jm + m^2} = \left(\frac{U}{s}\right)^k \frac{(2J)! \mathcal{X}_J}{k!(2J+k)!} . \quad (8)$$

On summing both sides over k , we find

$$\mathcal{X}_J^{-1} = (2J)! \left(\frac{U}{s}\right)^{-J} I_{2J} \left(2\sqrt{\frac{U}{s}}\right) \quad (9)$$

where $I_n(z)$ is the modified Bessel function of the first kind (ABRAMOWITZ and STEGUN, 1964). The fraction $\mathcal{X}_J(k)$ with $J+k$ mutations is then given by

$$\mathcal{X}_J(k) = \left(\frac{U}{s}\right)^{J+k} \frac{1}{k!(2J+k)!} \frac{1}{I_{2J} \left(2\sqrt{\frac{U}{s}}\right)} \quad (10)$$

and is plotted in Fig. 1a as function of $J+k$ at a fixed U/s . From the above equation, we find that for a given J , the fraction $\mathcal{X}_J(k)$ is centred about $k_2^* = \sqrt{J^2 + J_2^2} - J$ where $J_2 = \sqrt{U/s}$. For $J \ll J_2$, the distribution $\mathcal{X}_J(k)$ peaks about $k_2^* \approx J_2$ while for $J \gg J_2$, it is maximised at $J_2^2/(2J)$ (see Fig. 1a). Thus for large J , as argued at the beginning of this section, the distribution $\mathcal{X}_J(k)$ localises close to $k = 0$. The behavior of the least-loaded fraction $\mathcal{X}_J$ shown in Fig. 1b also depends on J_2 . For large J_2 (i.e. weak selection), the fraction $\mathcal{X}_J$ increases towards unity slower than for small J_2 . Using the asymptotic expansion of the Bessel function $I_n(z)$ for large orders (ABRAMOWITZ and STEGUN, 1964) in (9), we have

$$\mathcal{X}_J^{-1} \sim \frac{(2J)!}{\sqrt{4\pi J}} \left(\frac{U}{s}\right)^{-J} \frac{e^{2J\sqrt{1+y^2}}}{(1+y^2)^{1/4}} \left(\frac{y}{1+\sqrt{1+y^2}}\right)^{2J} \quad (11)$$




Figure 1: Steady state of the quasispecies model on synergistic fitness landscape with $\alpha = 2$: (a) Fraction $\chi_J(k)$ as a function of $J + k$ for $U/s = 50$ given by (10) (b) Fraction χ_J of the least-loaded class J calculated using (9).

where we have defined $y = J_2/J$. For $J \gg J_2$, the above expression can be simplified to give $\chi_J \approx \exp(-U/(2sJ))$ which asymptotically approaches unity. Thus with increasing J , most of the population tends to stay in the least-loaded class.

For arbitrary $\alpha > 1$, it does not seem possible to obtain explicit expression for $\chi_J(k)$. However using the insights obtained from $\alpha = 2$ case, we can find χ_J for large J . We expect that for any $\alpha > 1$, there exists a least-loaded class J_α such that the population frequency $\chi_J(k)$ with $J \gg J_\alpha$ is nonzero for $k \ll J$. In such a case, the denominator under the product sign in (6) can be expanded for $m \ll J$ to leading orders and yield

$$\chi_J(k) \approx \left(\frac{U}{\alpha s J^{\alpha-1}} \right)^k \frac{\chi_J}{k!}, \quad J \gg J_\alpha. \quad (12)$$

As $\chi_J(k)$ decays fast with k , we can sum over both sides of the above solution to obtain

$$\chi_J \approx \exp \left[-U/(\alpha s J^{\alpha-1}) \right], \quad J \gg J_\alpha. \quad (13)$$

This expression matches the exact results for $\alpha = 1$ and 2 discussed above. The product in (6) seems hard to calculate for $J \ll J_\alpha$. But for $J = 0$, we immediately have

$$\chi_0(k) \approx \left(\frac{U}{s} \right)^k \frac{\chi_0}{k!^\alpha} \quad (14)$$




Figure 2: Time evolution of the fraction $X(k, t)$ for $k = 12$ and 13 to illustrate the advance of the ratchet on epistatic fitness landscapes. Here $\alpha = 2$, $N = 256$, $U = 0.25$ and $s = 0.005$.

which peaks at k_α^* equal to $(U/s)^{1/\alpha}$. By analogy with the $\alpha = 2$ case, this suggests $J_\alpha = (U/s)^{1/\alpha}$.

TIME BETWEEN SUCCESSIVE CLICKS OF THE RATCHET

In this section, we first describe the process by which ratchet clicks and then calculate an expression for the typical time between successive clicks using a diffusion theory (EWENS, 1979). Let $X(k, t)$ denote the fraction of population with k mutations at time t in a single realization of the Wright-Fisher process. Figure 2 shows the time evolution of $X(k, t)$ for $k = 12$ and 13 starting from an initial condition in which all the N individuals carry zero mutations. As Fig. 2a illustrates, the fraction $X(12, t)$ increases from zero to a steady state fraction about which it fluctuates until a time $\tau_{11} = 1194$ after which it relaxes to another steady state before finally dropping to zero

at $\tau_{12} = 2130$ due to stochastic fluctuations. At $t = \tau_{12}$, the ratchet is said to have clicked as the least-loaded class with 12 mutations gets irreversibly lost and the class with 13 mutations shown in Fig. 2b becomes the new least-loaded class which itself gets lost at $\tau_{13} = 5235$. Since the ratchet is clicking at a slow rate, $X(k, t)$ has an opportunity to equilibrate. As Fig. 2b shows, soon after time τ_{12} , the fraction $X(13, t)$ fluctuates about a steady state fraction which is close to the deterministic frequency $\mathcal{X}_{13}$ given by (9). Similarly, $X(12, t)$ in Fig. 2a oscillates about $\mathcal{X}_{12}$ after the 11th error class is lost until time τ_{12} . As a click of the ratchet is signaled by the change in the average value of $X(k, t)$, it follows that the 11th class is lost at τ_{11} . For $\tau_{11} < t < \tau_{12}$ as there are at least 12 mutations in the population, the fraction $X(13, t)$ equilibrates about the frequency $\mathcal{X}_{12}(1)$. Thus the population fraction $X(k, t)$ for fixed k passes through a series of steady states with frequency $\mathcal{X}_J(k - J)$, $J \leq k$ before reaching the final absorbing state $X(k, \tau_k) = 0$. Note that this description of the mechanism by which the ratchet clicks assumes that the population $N\mathcal{X}_J$ in the currently least-loaded class J far exceeds one and thus has a chance to attain equilibrium before the next click.

We will use the diffusion approximation proposed in STEPHAN *et al.* (1993) to find the average inter-click time $\overline{T}_J = \overline{\tau_J - \tau_{J-1}}$ between the $(J-1)$ th and J th click of the ratchet where $\overline{}$ stands for averaging over stochastic histories. Let the random variable $X_J \in [0, 1]$ denote the population fraction in the least-loaded class J . If $X_J = 0$ at time t , the current least-loaded class J is lost forever and the ratchet is said to have clicked at t . We are interested in calculating the average time $\overline{T}_J$ required to reach the absorbing state $X_J = 0$ starting from X'_J at t' . The probability distribution $P(X_J, t|X'_J, t')$ obeys the following backward Fokker-Planck equation (RISKEN, 1996),

$$\begin{aligned}
-\frac{\partial P(X_J, t|X'_J, t')}{\partial t'} &= D_1(X'_J, t') \frac{\partial P(X_J, t|X'_J, t')}{\partial X'_J} \\
&+ \frac{D_2(X'_J, t')}{2} \frac{\partial^2 P(X_J, t|X'_J, t')}{\partial X'^2_J}
\end{aligned} \tag{15}$$

where

$$\begin{aligned}
D_n(X'_J, t') &= \text{Lt}_{\tau \rightarrow 0} \int_0^1 dX_J \frac{(X_J - X'_J)^n P(X_J, t' + \tau|X'_J, t')}{\tau} \\
&= \text{Lt}_{\tau \rightarrow 0} \frac{[X_J(t' + \tau) - X_J(t')]^n}{\tau}.
\end{aligned} \tag{16}$$

As the coefficients D_n in (15) are independent of t' (see below), the average inter-click time $\bar{T}_J$ defined as

$$\bar{T}(X'_J) = \int_{-\infty}^0 dt' (-t') P(0, 0 | X'_J, t') \quad (17)$$

obeys the following ordinary differential equation,

$$-1 = D_1(X'_J) \frac{d\bar{T}}{dX'_J} + \frac{D_2(X'_J)}{2} \frac{d^2\bar{T}}{dX'^2_J} . \quad (18)$$

Since $X'_J = 0$ is an absorbing state, the solution to the above equation is subjected to the boundary condition $\bar{T}(0) = 0$. Furthermore as the population in the J th class equilibrates about the mean $\mathcal{X}_J$ after the $(J-1)$ th click, we can choose the initial distribution of random variable X'_J to be $\delta(X'_J - \mathcal{X}_J)$. Then the time $\bar{T}_J$ during which J is the least loaded class obtained by solving (18) is given by (EWENS, 1979)

$$\bar{T}_J = 2 \int_0^{\mathcal{X}_J} dY \psi(Y) \int_Y^1 \frac{dZ}{\psi(Z) D_2(Z)} \quad (19)$$

where $\psi(Y) = \exp \left[-2 \int^Y dX D_1(X) / D_2(X) \right]$.

We will now determine the coefficients D_1 and D_2 . The drift coefficient D_1 defined in (16) measures the change in the average fraction of the least-loaded class over a generation. As the population is in local equilibrium, this can be determined using the quasispecies equation (3) for $j = 0$. Thus the drift coefficient is given by (STEPHAN *et al.*, 1993; GORDO and CHARLESWORTH, 2000a)

$$\begin{aligned} D_1(X_J = \mathcal{X}_J(0, t)) &= \mathcal{X}_J(0, t+1) - \mathcal{X}_J(0, t) \\ &= X_J \frac{e^{-U} W(J) - \mathcal{W}_J(t)}{\mathcal{W}_J(t)} . \end{aligned} \quad (20)$$

As expected, D_1 vanishes when the population is either in the steady state (see (5)) or in the absorbing state ($X_J = 0$). The equation (20) for D_1 does not close in X_J but one can obtain an approximate expression for D_1 using the linear response theory (RISKEN, 1996). As D_1 is proportional to the deviation from a steady state quantity, we can write (STEPHAN *et al.*, 1993)

$$D_1 \propto e^{-U} W(J) - \mathcal{W}_J(t) = C \left(1 - \frac{X_J}{\mathcal{X}_J} \right) \quad (21)$$

where C is a constant. Thus the drift coefficient can be written in terms of X_J as

$$D_1 \approx CX_J \left(1 - \frac{X_J}{\mathcal{X}_J}\right) . \quad (22)$$

As Fig. 2 shows, when the ratchet clicks ($X_J = 0$) and the J th class is lost, the population quickly relaxes to the equilibrium frequency of the $(J+1)$ th class so that the deviation in the fitness $C \sim \mathcal{W}_J - \mathcal{W}_{J+1} = e^{-U} [W(J) - W(J+1)]$ where we have used (5). For $s \rightarrow 0$, expanding the fitness $W(J)$ to leading orders in s , we get

$$D_1(X_J) \approx AX_J \left(1 - \frac{X_J}{\mathcal{X}_J}\right) \quad (23)$$

where $A = cs [(J+1)^\alpha - J^\alpha] \approx \alpha cs J^{\alpha-1}$ for large J .

The diffusion coefficient D_2 in (16) gives the fluctuations in the frequency of the least-loaded class about the mean value. These fluctuations arising due to the finiteness of the population can be determined using (2) which gives the variance in the number of offspring produced in one generation as (EWENS, 1979)

$$D_2(X_J) = \frac{X_J(1 - X_J)}{N} \approx \frac{X_J}{N} . \quad (24)$$

The last expression on the right hand side of the above equation captures the fact that the fluctuations vanish when either the population size N is infinite or the population is in the absorbing state $X_J = 0$.

Using the coefficients (23) and (24) in (19), the average inter-click time $\bar{T}_J$ can be written as

$$\bar{T}_J = 2N \int_0^{\mathcal{X}_J} dY \psi(Y) \int_Y^1 \frac{dZ}{Z} \psi^{-1}(Z) \quad (25)$$

where

$$\psi(Y) = \exp \left[\beta \left(\frac{Y^2}{\mathcal{X}_J^2} - \frac{2Y}{\mathcal{X}_J} \right) \right] \quad (26)$$

and

$$\beta = N\mathcal{X}_J A = N\mathcal{X}_J cs [(J+1)^\alpha - J^\alpha] . \quad (27)$$

In the absence of epistasis, both A and $\mathcal{X}_J$ are independent of J so that typical time spent between any two successive clicks is constant and the ratchet turns with a finite speed equal to $1/\bar{T}$. For epistatic fitness $\alpha \neq 1$, $\bar{T}_J$ depends explicitly on J and is expected to increase with J for $\alpha > 1$

while decrease for $\alpha < 1$ (KONDRASHOV, 1994). In the following discussion, we will restrict ourselves to $\alpha \geq 1$.

After some simple manipulations, we can rewrite (25) as

$$\bar{T}_J = 2N\mathcal{X}_J \int_{-1}^0 dY e^{\beta Y^2} \int_Y^\delta \frac{dZ}{1+Z} e^{-\beta Z^2} \quad (28)$$

which implies that the scaled time $\bar{T}_J/(N\mathcal{X}_J)$ is a function of two variables namely $\beta = N\mathcal{X}_J A$ and $\delta = (1 - \mathcal{X}_J)/\mathcal{X}_J$. The nature of $\bar{T}_J$ depends on the parameter $\beta\delta^2$ which can be seen as follows. Consider the Gaussian $e^{-\beta Z^2}$ in the rightmost integral in (28) which is centred about $Z = 0$ and has a width $1/\sqrt{\beta}$. If the upper limit δ of this integral exceeds the width i.e. $\beta\delta^2 \gg 1$, the integral can be cutoff at $1/\sqrt{\beta}$ thus eliminating the dependence on δ . In such a case, $\bar{T}_J$ is of the following scaling form,

$$\bar{T}_J \approx N\mathcal{X}_J \mathcal{F}(N\mathcal{X}_J A) \quad (29)$$

where $\mathcal{F}(\beta)$ is the scaling function determined below. If $\beta\delta^2 \ll 1$, then the inter-click time depends on δ as well. Since $\beta\delta^2 \sim NsJ^{\alpha-1}/\mathcal{X}_J$ and $\mathcal{X}_J$ is bounded above by one, $\beta\delta^2 \gg 1$ for large J when epistatic interactions are synergistic. For $\alpha = 1$, the parameter $\beta\delta^2$ exceeds unity if $Ns \gg 1$. Here we will restrict ourselves to the $\beta\delta^2 \gg 1$ case which has nice scaling properties although the double integral in (28) can be estimated for $\beta\delta^2 \ll 1$ also.

We now proceed to find the scaling function $\mathcal{F}(\beta)$. If the width of the Gaussian is large i.e. $\beta \ll 1$, we can approximate $e^{-\beta Z^2} \approx 1$ for $Z \ll 1/\sqrt{\beta}$ in the rightmost integral in (28). Since $\delta \gg 1/\sqrt{\beta}$, as argued above, this integral needs to be carried out from Y to $1/\sqrt{\beta}$. This yields

$$\begin{aligned} \bar{T}_J &\approx 2N\mathcal{X}_J \int_0^1 dY e^{\beta Y^2} \left[\ln \left(1 + \frac{1}{\sqrt{\beta}} \right) - \ln(1 - Y) \right] \\ &\approx 2N\mathcal{X}_J \left[1 + \ln \left(1 + \frac{1}{\sqrt{\beta}} \right) \right] \\ &\approx N\mathcal{X}_J (2 - \ln \beta) , \quad \beta \ll 1 . \end{aligned} \quad (30)$$

To find the scaling function in the opposite limit $\beta \gg 1$, we first consider the inner integral in (28),

$$\begin{aligned} \int_Y^\delta \frac{dZ}{1+Z} e^{-\beta Z^2} &= \frac{1}{\sqrt{\beta}} \int_{Y\sqrt{\beta}}^{\delta\sqrt{\beta}} dZ e^{-Z^2} \left[1 + \frac{Z}{\sqrt{\beta}} \right]^{-1} \\ &\approx \frac{1}{2} \sqrt{\frac{\pi}{\beta}} \left[\text{erf}(\delta\sqrt{\beta}) - \text{erf}(Y\sqrt{\beta}) \right] \end{aligned} \quad (31)$$

where $\text{erf}(z)$ is the error function (ABRAMOWITZ and STEGUN, 1964) and we have kept terms to leading orders in $\beta\delta^2(\gg 1)$ and $Y\sqrt{\beta}$ for $\beta \gg 1$. On using the above integral in (28), we obtain

$$\begin{aligned}\bar{T}_J &\approx \frac{N\mathcal{X}_J\sqrt{\pi}}{\beta} \int_0^{\sqrt{\beta}} dY e^{Y^2} \left[\text{erf}(\delta\sqrt{\beta}) + \text{erf}(Y) \right] \\ &= \frac{N\mathcal{X}_J\sqrt{\pi}e^\beta}{2\beta} \int_0^\beta \frac{dY e^{-Y}}{\sqrt{\beta-Y}} \left[\text{erf}(\delta\sqrt{\beta}) + \text{erf}(\sqrt{\beta-Y}) \right] \\ &\approx N\mathcal{X}_J \sqrt{\frac{\pi}{\beta}} \frac{e^\beta}{\beta}\end{aligned}\tag{32}$$

where we have used that both $\beta\delta^2$ and β are large. From the above discussion, we find that the scaling function

$$\mathcal{F}(\beta) \approx \begin{cases} 2 - \ln \beta, & \beta \ll 1 \\ \sqrt{\pi} \beta^{-3/2} e^\beta, & \beta \gg 1 \end{cases}\tag{33}$$

is a U-shaped function of β reaching a minimum when $\beta \sim 1$. We now discuss the above results in more detail for $\alpha = 1$ and $\alpha > 1$.

Non-epistatic fitness landscapes: As we have already argued, when $\alpha = 1$ the ratchet clicks with a finite speed equal to $1/\bar{T}$. When the rate at which the ratchet clicks is large, analytical results can be obtained using the traveling wave approach (ROUZINE *et al.*, 2003). For slowly clicking ratchet which is the subject of this article, the problem was formulated analytically within a diffusion approximation first by STEPHAN *et al.* (1993). However a better agreement between the diffusion theory and the simulation results was obtained in GORDO and CHARLESWORTH (2000a,b). A possible reason for this difference is that STEPHAN *et al.* (1993) included terms besides those in (22) in the expansion of the drift coefficient which is not consistent with the assumption of linear response, while the expression for D_1 in GORDO and CHARLESWORTH (2000a) is the same as (22). In fact, the expression for the inter-click time given by (28) with $c = 0.6$ is identical to that reported in GORDO and CHARLESWORTH (2000a). However the integrals were computed numerically by these authors while here we estimate them analytically and find that the average inter-click time is given by

$$\bar{T} \approx \begin{cases} n_0 (2 - \ln \beta), & \beta \ll 1 \\ \sqrt{\pi} n_0 \beta^{-3/2} e^\beta, & \beta \gg 1 \end{cases}\tag{34}$$




Figure 3: Data collapse of the scaled time $\bar{T}/n_0$ when plotted against the scaling parameter $\beta = n_0cs$ for the multiplicative fitness landscape where $c = 0.6$. The simulation data has been averaged over 100 stochastic runs.

where $n_0 = N\mathcal{X}_J = Ne^{-U/s}$ is the number of individuals in the least-loaded class and $\beta = n_0cs$ is the scaling parameter. The inter-click time calculated numerically using the integral (28) and the above expression (34) is shown in Table 1 and the two are seen to be in good agreement.

Our first result concerning the Muller's ratchet is the scaling form for time $\bar{T}$ when parameters N, U and s are chosen such that $\beta\delta^2 \sim Ns \gg 1$ is satisfied. The results of our numerical simulations testing this scaling form are shown in Fig. 3 where we have differentiated data points at fixed β for clarity. The scaled time $\bar{T}/n_0$ indeed shows a very good data collapse and a non-monotonic dependence on β . HIGGS and WOODCOCK (1995) unsuccessfully attempted to obtain a scaling form for the ratchet rate by using Ns as the scaling parameter for $Ns \gg 1$. The scaling function in (34) however shows that the scaling parameter is a function of the population number n_0 in the least-loaded class, and not the total population N .

In many studies (HAIGH, 1978; BELL, 1988; GESSLER, 1995), the size n_0 of the least-loaded class has been regarded as an important parameter in

β	N	U	s	n_0	$\overline{T}_{\text{sim}}(2 \text{ SE})$	$\overline{T}_{\text{int}}$	$\overline{T}_{\text{sca}}$
0.121	7500	0.02	0.004	50.5	453(34)	230	207
	15000	0.01	0.002	101.1	906(69)	460	416
	30000	0.005	0.001	202.1	1805(111)	920	831
0.242	3000	0.1	0.02	20.2	141(7)	86	69
	6000	0.05	0.01	40.4	271(23)	172	138
	12000	0.025	0.005	80.8	559(39)	344	276
1.212	15000	0.1	0.02	101.1	506(41)	479	451
	30000	0.05	0.01	202.2	1008(101)	959	902
	60000	0.025	0.005	404.2	1950(133)	1918	1804
2.424	30000	0.1	0.02	202.1	1490(149)	1487	1071
	45000	0.067	0.013	303.2	2104(218)	2230	1608
	60000	0.05	0.01	404.2	2579(257)	2974	2143

Table 1: Comparison of simulation data plotted in Fig. 3 with analytical results. Here $\overline{T}_{\text{sim}}$ refers to the inter-click time obtained in simulations, $\overline{T}_{\text{int}}$ by numerically evaluating the integrals in (28) for $\alpha = 1$ and $\overline{T}_{\text{sca}}$ using the scaling form (34).

determining the ratchet speed. If $n_0 \gg 1$, the population is close to the deterministic limit and the ratchet clicks slowly whereas for $n_0 \ll 1$, the ratchet speed is high. However the simulations show that the size n_0 of the least-loaded class is not sufficient to predict the ratchet rate (STEPHAN *et al.*, 1993; GORDO and CHARLESWORTH, 2000a). This is indeed captured by diffusion approximation as (34) is not a function of n_0 alone. However, if s is kept fixed, $\overline{T}$ increases monotonically with n_0 in accordance with the above expectation and simulations (GABRIEL *et al.*, 1993; GORDO and CHARLESWORTH, 2000a). The dependence of $\overline{T}$ on s for given n_0 is however non-monotonic similar to that seen in numerical studies (GORDO and CHARLESWORTH, 2000a). To understand this behavior qualitatively, consider the situation where the population number n_0 is kept constant by keeping N and U/s fixed. As increasing s tends to localise population and hence increase $\overline{T}$, increasing U has the opposite delocalising effect which decreases $\overline{T}$. At a given U and s , one of these two competing forces wins. According to (34), as the scaling function overturns when $n_0 s \sim 1$, the mutation takes over for $U > U^* = s \ln(Ns)$ whereas below U^* , selection dominates and $\overline{T}$ is large.

The solution (34) gives an initial logarithmically slow drop in s and an exponential increase for larger s with the minimum of the U-shaped curve occurring at a selection coefficient which scales as $1/n_0$. The simulation results of GORDO and CHARLESWORTH (2000a) (also see Table 1) however show a much faster drop at small s . A good agreement with simulation data was obtained in GORDO and CHARLESWORTH (2000b) by adding $\bar{T}$ and time $T_a \sim 1/s$ required by the population to relax to new steady state just after a click. However, the full expression for T_a is not of scaling form (29) although the simulation data in Fig. 3 shows an excellent data collapse even for small s . In view of this, a better understanding of the time T_a is desirable. Of course, for both n_0 and s fixed, the time $\bar{T}$ is predicted to be independent of N and U which is confirmed by simulations (GORDO and CHARLESWORTH, 2000a).

Epistatic fitness landscapes: For synergistic interactions, the ratchet is expected to halt at large times (CHARLESWORTH *et al.*, 1993; KONDRASHOV, 1994). Figure 4a shows the population fraction $\bar{X}(k, t)$ with k deleterious mutations at several time slices. Two points are noteworthy: the ratchet does not turn with a constant speed as is evident by the rate at which the population accumulates average number of mutations at late times. Secondly, unlike for the multiplicative fitness case (ROUZINE *et al.*, 2003), the population fraction does not maintain its shape as the width of the distribution $\bar{X}(k, t)$ decreases with increasing time. Thus the number frequency of an asexual population under epistatic selection does not behave like a traveling wave moving with a constant speed.

The simulation data for average inter-click time shown in Fig. 4b increases with the minimum number J of deleterious mutations in the population thus indicating the arrest of the ratchet at large times. The time $\bar{T}_J$ obtained by integrating (28) for $\alpha = 2$ is also shown for comparison and we find that it agrees well with the simulation results. As the scaling parameter β defined in (27) increases as a power law with J for $J \gg 1$, we have $\beta \gg 1$ for large J . In such a case, $\bar{T}_J \sim N\mathcal{X}_J\beta^{-3/2}e^\beta$ which increases exponentially fast with β . As $\mathcal{X}_J \rightarrow 1$ for large J , the inter-click time $\bar{T}_J \sim e^{N\alpha csJ^{\alpha-1}}$ increases faster than any power law with J and for any $\alpha > 1$. Thus an arbitrarily small $\alpha - 1$ is capable of slowing down the ratchet under synergistic selection.

To estimate how the ratchet speed approaches zero, we use the following argument. The average speed $\bar{v} = d/t$ where d is the number of minimum mutations accumulated in time interval t . Since T_J is the time between the




Figure 4: Loss of least-loaded class on epistatic fitness landscapes with $\alpha = 2$, $N = 256$, $U = 0.25$, $s = 0.005$: (a) Average fraction $\bar{X}(k, t)$ in the k th class at $t = 50, 100, 200, 1250, 2500, 5000, 10000$ (from left to right). The data have been averaged over 1000 stochastic histories. (b) Average inter-click time $\bar{T}_J$ as a function of J where each data point has been averaged over 500 histories. Comparison with $\bar{T}_J$ obtained by evaluating integral (28) numerically and using the scaling form (29) at large J is also shown. A reasonably good agreement is obtained with $c = 0.9$.

$(J - 1)$ th and J th click of the ratchet, we have

$$\bar{v} = \frac{d}{T_0 + \dots + T_d} \quad (35)$$

where d is fixed. Assuming the distribution for click times has nice scaling properties, we may write

$$\bar{v} \sim \frac{d}{\overline{T_0 + \dots + T_d}}. \quad (36)$$

If T_J does not depend on the least-loaded class J , we have $\bar{v} = 1/\bar{T}$ as expected for $\alpha = 1$ case. For $\alpha > 1$ as the average inter-click time increases faster than power law with J , the sum $\overline{T_0 + \dots + T_d} \sim d^{2-\alpha} e^{N\alpha cs d^{\alpha-1}}$ which is reasonable as the sum is dominated by $\overline{T_d}$ for large d . This gives $\bar{v} \sim d^{\alpha-1} e^{-N\alpha cs d^{\alpha-1}}$ or in terms of t ,

$$\bar{v} \sim \frac{1}{t} \left(\frac{\ln t}{N\alpha cs} \right)^{1/(\alpha-1)}. \quad (37)$$

Thus in the presence of epistasis, the average speed of the ratchet approaches zero as $1/t$ with α -dependent logarithmic corrections.

DISCUSSION

In this article, we considered the effect of drift and epistasis on the loss of the least-loaded (or the fittest) class in an asexual population. When the population size is infinite, the drift is absent and the population evolves due to the elementary processes of selection and mutation. As selection tends to localise the population at the fittest sequence while mutation has the opposite tendency to delocalise it, an error threshold may exist beyond which the fittest class cannot be sustained in the population (EIGEN, 1971). Such a phase transition is known to occur for asexual populations evolving deterministically on fitness landscapes defined by (1) when $\alpha < 1$ (WIEHE, 1997). However for $\alpha \geq 1$, the population frequency $\mathcal{X}_J$ in the least-loaded class J remains nonzero for any finite U, s . In fact, for synergistic interactions, the frequency $\mathcal{X}_J$ given by (13) increases with J towards unity.

If however the population size is finite, as illustrated in Fig. 2, the population frequency $X_J(t)$ fluctuates with time and can become zero even for $\alpha \geq 1$. Numerical simulations have shown that this loss occurs at a constant

speed when $\alpha = 1$ (HAIGH, 1978) but at a decelerating rate when $\alpha > 1$ (CHARLESWORTH *et al.*, 1993; KONDRASHOV, 1994). In this paper, we focused on the stochastic dynamics of the loss of the fittest class for $\alpha \geq 1$ and calculated explicit analytical expressions for the typical time $\bar{T}_J$ during which the least-loaded class J survives using a diffusion theory (EWENS, 1979). Although this approach has been considered previously to attack the Muller’s ratchet problem (STEPHAN *et al.*, 1993), the resulting solution in the form of a double integral was evaluated numerically which does not allow one to infer the functional dependence of $\bar{T}_J$ on parameters N, U, s and J .

When the interactions between gene loci are assumed to be absent and $Ns \gg 1$, the inter-click time $\bar{T}$ is found to be of the scaling form (34). Although this result is derived using diffusion theory which is based on several approximations, the numerical simulations show an excellent data collapse suggesting that (29) may be an exact statement. For fixed n_0 , the time $\bar{T}$ is seen to be a U-shaped function of s arising due to competition between mutation and selection. Such a behavior is reminiscent of error threshold phenomenon in infinite populations discussed above. Although the least-loaded class is never lost in the deterministic limit on multiplicative fitness landscapes (WAGNER and KRALL, 1993), the selection-mutation competition manifests itself in the time duration during which the least-loaded (fittest) class can support a finite population. For given s , the survival time $\bar{T}$ initially increases linearly with n_0 approaching the deterministic limit of $N \rightarrow \infty$ with an exponential rise with n_0 as increasing N decreases the effect of drift.

Muller’s ratchet has been proposed as a possible mechanism for the degeneration and eventual extinction of asexual organisms and nonrecombining parts of sexually reproducing populations. For bacterial populations with $N \sim 10^6$, $U \approx 0.003$ (DRAKE *et al.*, 1998) and $s \approx 10^{-3}$, using (34) we find it takes 10^{15} generations for one click to occur which does not seem plausible. However, reducing s by a factor half gives $\bar{T} \sim 5000$ and further reduction to $s \approx 0.25 \times 10^{-3}$ gives just 50 generations for a single click. Muller’s ratchet has also been invoked to understand the degeneration of the nonrecombining neo-Y chromosome which originated about a million years ago in *Drosophila Miranda* (CHARLESWORTH, 1978; GORDO and CHARLESWORTH, 2000a). Assuming that a few thousand deleterious mutations have occurred over this time span, the time between successive turns of the ratchet is of the order of 10^3 generations. From (34), the time $\bar{T} \sim e^{n_0 cs}$ for large $\bar{T}$ which gives $n_0 s \approx 14$. If $s \approx 10^{-2}$, the population size N required for the

Muller’s ratchet to be a viable mechanism works out to be $\sim 700e^{100U}$ which depends sensitively on U due to the exponential dependence. For instance, for $U = 0.07$, the required population is of the order 5×10^5 while it reduces by a factor twenty for $U = 0.04$. A similar sensitive dependence of extinction time on s for given N and U has been noted in the problem of the degeneration of human mitochondrial DNA also (LOEWE, 2006). This suggests that very precise estimates of U and s may be required to determine whether the Muller’s ratchet might be in operation.

The scaling form (29) holds for $\alpha > 1$ also but the speed of the ratchet under epistatic selection is found to decay rapidly with time for any $\alpha > 1$. Although the Muller’s ratchet under synergistically epistatic selection has the interesting feature of arresting the loss of least-loaded class, the generality of this mechanism seems unclear as general support for synergistic epistasis has not been found in the experiments (DE VISSER and ELENA, 2007) and the slowing down effect is sensitive to the inclusion of biologically relevant details such as distribution of mutational effects (BUTCHER, 1995). Recent experimental evidence suggests that fitness decline down to a plateau can be attributed to the presence of epistasis (SILANDER *et al.*, 2007). This can be due to negative epistasis (KONDRASHOV, 1994) or compensatory epistasis (SILANDER *et al.*, 2007). It would be interesting to compare the results discussed in this work with models that include compensatory mutations (WAGNER and GABRIEL, 1990; WILKE *et al.*, 2003).

Acknowledgement: The author is grateful to J. Krug, S.-C. Park and an anonymous referee for helpful comments on the manuscript.

References

- ABRAMOWITZ, M. and I. STEGUN, 1964 *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*. Dover.
- BACHTROG, D. and I. GORDO, 2004 Adaptive evolution of asexual populations under Muller’s ratchet. *Evolution* **58**: 1403–1413.
- BARTON, N. and B. CHARLESWORTH, 1998 Why sex and recombination? *Science* **281**: 1986–1990.
- BELL, G., 1988 Recombination and the immortality of the germ line. *J. evol. Biol.* **1**: 67–82.

- BUTCHER, D., 1995 Muller's ratchet, epistasis and mutation effects. *Genetics* **141**: 431–437.
- CHARLESWORTH, B., 1978 Model for evolution of Y chromosomes and dosage compensation. *Proc. Natl. Acad. Sci. USA* **75**: 5618–5622.
- CHARLESWORTH, D., M. MORGAN, and B. CHARLESWORTH, 1993 Mutation accumulation in finite outbreeding and inbreeding populations. *Genet. Res. Camb.* **61**: 39–56.
- COMBADÃO, J., P. CAMPOS, F. DIONISIO, and I. GORDO, 2007 Small-world networks decrease the speed of Muller's ratchet. *Genet. Res.* **89**: 7–18.
- DE VISSER, J. and S. ELENA, 2007 The evolution of sex: empirical insights into the roles of epistasis and drift. *Nat. Rev. Genet.* **8**: 139–149.
- DRAKE, J.W., B. CHARLESWORTH, D. CHARLESWORTH, and J.F. CROW, 1998 Rates of spontaneous mutation. *Genetics* **148**: 1667–1686.
- EIGEN, M., 1971 Selforganization of matter and evolution of biological macromolecules. *Naturwissenschaften* **58**: 465 – 523.
- EWENS, W., 1979 *Mathematical Population Genetics*. Springer, Berlin.
- FELSENSTEIN, J., 1974 The evolutionary advantage of recombination. *Genetics* **78**: 737–756.
- GABRIEL, W., M. LYNCH, and R. BÜRGER, 1993 Muller's ratchet and mutational meltdowns. *Evolution* **47**: 1744–1757.
- GESSLER, D. D. G., 1995 The constraints of finite size in asexual populations and the rate of the ratchet. *Genetical Research* **66**: 241–253.
- GORDO, I. and B. CHARLESWORTH, 2000a The degeneration of asexual haploid populations and the speed of Muller's ratchet. *Genetics* **154**: 1379–1387.
- GORDO, I. and B. CHARLESWORTH, 2000b On the speed of Muller's ratchet. *Genetics* **156**: 2137–2140.

- HAIGH, J., 1978 The accumulation of deleterious genes in a population - Muller's ratchet. *Theoret. Population Biol.* **14**: 251–267.
- HIGGS, P., 1994 Error thresholds and stationary mutant distributions in multi-locus diploid genetics models. *Genet. Res. Camb.* **63**: 63–78.
- HIGGS, P. G. and G. WOODCOCK, 1995 The accumulation of mutations in asexual populations and the structure of genealogical trees in the presence of selection. *J. Math. Biol.* **33**: 677–702.
- HURST, L. and J. PECK, 1996 Recent advances in understanding of the evolution and maintenance of sex. *Trends Ecol. Evol.* **11**: 46–52.
- JAIN, K. and J. KRUG, 2007 Adaptation in simple and complex fitness landscapes. In U. Bastolla, M. Porto, H. Roman, and M. Vendruscolo (Eds.), *Structural Approaches to Sequence Evolution: Molecules, Networks and Populations*, pp. 299–340. Springer, Berlin.
- JUDSON, O. and B. NORMARK, 1996 Ancient asexual scandals. *Trends Ecol. Evol.* **11**: 41–46.
- KIMURA, M. and T. MARUYAMA, 1966 The mutational load with epistatic gene interactions in fitness. *Genetics* **54**: 1337–1351.
- KONDRASHOV, A., 1994 Muller's ratchet under epistatic selection. *Genetics* **136**: 1469–1473.
- LOEWE, L., 2006 Quantifying the genomic decay paradox due to Muller's ratchet in human mitochondrial DNA. *Genet. Res. Camb.* **87**: 133–159.
- MULLER, H. J., 1964 The relation of recombination to mutational advance. *Mutation Res.* **1**: 2–9.
- PAMILO, P., M. NEI, and W.-H. LI, 1987 Accumulation of mutations in sexual and asexual populations. *Genet. Res.* **49**: 135–146.
- PRÜGEL-BENNETT, A., 1997 Modelling evolving populations. *J. theo. Biol.* **185**: 81–95.
- RISKEN, H., 1996 *The Fokker Planck equation. Methods of solution and applications*. Springer, Berlin.

- ROUZINE, I., J. WAKELEY, and J. COFFIN, 2003 The solitary wave of asexual evolution. *Proc. Natl. Acad. Sci. U.S.A.* **100**: 587–592.
- SCHULTZ, S. and M. LYNCH, 1997 Mutation and extinction: the role of variable mutational effects, synergistic epistasis, beneficial mutations, and degree of outcrossing. *Evolution* **51**: 1363–1371.
- SILANDER, O., O. TENAILLON, and L. CHAO, 2007 Understanding the Evolutionary Fate of Finite Populations: The Dynamics of Mutational Effects. *PLoS Biology* **5**: 922–931.
- STEPHAN, W., L. CHAO, and J. G. SMALE, 1993 The advance of Muller ratchet in a haploid asexual population - approximate solutions based on diffusion-theory. *Genet. Res. Camb.* **61**: 225–232.
- STEPHAN, W. and Y. KIM, 2007 Recent application of diffusion theory to population genetics. In M. Slatkin and M. Veuille(Eds.), *Modern Developments in Theoretical Population Genetics*, pp. 72–93. Oxford University Press, Oxford.
- WAGNER, G. and W. GABRIEL, 1990 Quantitative Variation in Finite Parthenogenetic Populations: What Stops Muller’s Ratchet in the Absence of Recombination? *Evolution* **44**: 715–731.
- WAGNER, G. and P. KRALL, 1993 What is the difference between models of error thresholds and Muller’s ratchet? *J. Math. Biol.* **32**: 33–44.
- WIEHE, T., 1997 Model dependency of error thresholds: the role of fitness functions and contrasts between the finite and infinite sites models. *Genet. Res. Camb.* **69**: 127–136.
- WILKE, C., R. LENSKE, and C. ADAMI, 2003 Compensatory mutations cause excess of antagonistic epistasis in RNA secondary structure folding. *BMC Evol. Biol.* **3**: 3.
- WOLF, J., E. BRODIE, and M. WADE (Eds.), 2000 *Epistasis and the evolutionary process*. Oxford Univ. Press, New York.
- WOODCOCK, G. and P. G. HIGGS, 1996 Population evolution on a multiplicative single-peak fitness landscape. *J. theor. Biol.* **179**: 61–73.